

L-S CATEGORIES OF SIMPLY-CONNECTED COMPACT SIMPLE LIE GROUPS OF LOW RANK

NORIO IWASE AND MAMORU MIMURA

ABSTRACT. We determine the L-S category of $Sp(3)$ by showing that the 5-fold reduced diagonal $\overline{\Delta}_5$ is given by ν^2 , using a Toda bracket and a generalised cohomology theory h^* given by $h^*(X, A) = \{X/A, \mathbb{S}[0, 2]\}$, where $\mathbb{S}[0, 2]$ is the 3-stage Postnikov piece of the sphere spectrum $\mathbb{S}$. This method also yields a general result that $\text{cat}(Sp(n)) \geq n + 2$ for $n \geq 3$, which improves the result of Singhof [17].

1. INTRODUCTION

IN THIS paper, each space is assumed to have the homotopy type of a CW complex. The (normalised) L-S category of X is the least number m such that there is a covering of X by $(m + 1)$ open subsets each of which is contractible in X . Hence $\text{cat}\{*\} = 0$. By Lusternik and Schnirelmann [10], the number of critical points of a smooth function on a manifold M is bounded below by $\text{cat } M + 1$.

G. Whitehead showed that $\text{cat}(X)$ coincides with the least number m such that the diagonal map $\Delta_{m+1} : X \rightarrow \prod^{m+1} X$ can be compressed into the ‘fat wedge’ $T^{m+1}(X)$ (see Chapter X of [20]). Since $\prod^{m+1} X / T^{m+1}(X)$ is the $(m + 1)$ -fold smash product $\wedge^{m+1} X$, we have a weaker invariant $w\text{cat } X$, the *weak L-S category* of X , given by the least number m such that the reduced diagonal map $\overline{\Delta}_{m+1} : X \rightarrow \wedge^{m+1} X$ is trivial. Hence $w\text{cat } X \leq \text{cat } X$.

T. Ganea has also introduced a stronger invariant $\text{Cat } X$, the *strong L-S category* of X , by the least number m such that there is a covering of X by $(m + 1)$ open subsets each of which is contractible in itself. Thus $w\text{cat } X \leq \text{cat } X \leq \text{Cat } X$.

The weak and strong L-S categories usually give nice estimates of L-S category especially for manifolds. Actually, we do not know any example of a closed manifold whose strong L-S, L-S and weak L-S categories are not the same. The following problems are posed by Ganea [4]:

- i) (Problem 1) Determine the L-S category of a manifold.
- ii) (Problem 4) Describe the L-S category of a sphere-bundle over a sphere in terms of homotopy invariants of the characteristic map of the bundle.

Problem 1 has been studied by many authors, such as Singhof [16, 17, 18], Montejano [12], Schweizer [15], Gomez-Larrañaga and Gonzalez-Acuña [5], James and Singhof [9] and Rudyak [13, 14]. In particular for compact simply-connected simple Lie groups, $\text{cat}(SU(n + 1)) = n$ for $n \geq 1$ by [16], $\text{cat}(Sp(2)) = 3$ by [15] and

Date: December 17th, 2001, third draft.

1991 Mathematics Subject Classification. Primary 55M30, Secondary 55N20, 22E20.

Key words and phrases. Lie group, Lusternik-Schnirelmann category, cohomology theory.

The second named author is partially supported by the Grant-in-Aid for Scientific Research #12640025 from the Japan Society of Promotion of Science.

$\text{cat}(Sp(n)) \geq n + 1$ for $n \geq 2$ by [17]. It was also announced recently that Problem 4 was solved by the first author [7].

The method in the present paper also provides a result for G_2 , and thus we have the following result.

Theorem 1.1. *The following is the complete list of L-S categories of a simply-connected compact simple Lie group of rank ≤ 2 :*

Lie groups	$Sp(1) = SU(2) = Spin(3)$	$SU(3)$	$Sp(2) = Spin(5)$	G_2
wcat	1	2	3	4
cat	1	2	3	4
Cat	1	2	3	4

Although the above result is known for experts, we give a short proof for G_2 . In fact, the result for G_2 has never been published and is obtained in a similar but easier manner than the following result for $Sp(3)$:

Theorem 1.2. $wcat(Sp(3)) = \text{cat}(Sp(3)) = \text{Cat}(Sp(3)) = 5$.

Remark 1.3. *The argument given to prove Theorem 1.2 provides an alternative proof of Schweizer's result*

$$wcat(Sp(2)) = \text{cat}(Sp(2)) = \text{Cat}(Sp(2)) = 3.$$

The authors know that a similar result to Theorem 1.2 is obtained by Lucía Fernández-Suárez, Antonio Gómez-Tato, Jeffrey Strom and Daniel Tanré [3]. Our method is, however, much simpler and providing the following general result:

Theorem 1.4. $n + 2 \leq wcat(Sp(n)) \leq \text{cat}(Sp(n)) \leq \text{Cat}(Sp(n))$ for $n \geq 3$.

This improves Singhof's result: $\text{cat}(Sp(n)) \geq n + 1$ for $n \geq 2$. We propose the following conjecture.

Conjecture 1.5. *Let G be a simply-connected compact Lie group with $G = \prod_{i=1}^n H_i$ where H_i is a simple Lie group. Then $wcat(G) = \text{cat}(G) = \text{Cat}(G)$ and $\text{cat}(G) = \sum_{i=1}^n \text{cat}(H_i)$.*

It might be difficult to say something about $\text{cat } Sp(n)$, but an old conjecture says the following.

Conjecture 1.6. $\text{cat } Sp(n) = 2n - 1$ for all $n \geq 1$.

The authors thank John Harper for many helpful conversations.

2. PROOF OF THEOREM 1.1

Let us recall a CW decomposition of G_2 from [11]:

$$G_2 = e^0 \cup e^3 \cup e^5 \cup e^6 \cup e^8 \cup e^9 \cup e^{11} \cup e^{14}.$$

On the other hand, we have the following cone-decomposition.

Theorem 2.1. *There is a cone-decomposition of G_2 as follows:*

$$\begin{aligned} G_2^{(5)} = \Sigma \mathbb{C}P^2, \quad S^5 \cup e^7 \rightarrow G_2^{(5)} \hookrightarrow G_2^{(8)}, \\ S^8 \cup e^{10} \rightarrow G_2^{(8)} \hookrightarrow G_2^{(11)}, \quad S^{13} \rightarrow G_2^{(11)} \hookrightarrow G_2. \end{aligned}$$

Proof. The first and the last formulae are obvious. So we show the 2nd and 3rd formulae: By taking the homotopy fibre F_1 of $G_2^{(5)} \hookrightarrow G_2$, we can easily observe using the Serre spectral sequence that the fibre has a CW structure given by $S^5 \cup e^7 \cup (\text{cells in dimensions } \geq 7)$, where the cohomology generators corresponding to S^5 and e^7 are transgressive. Thus the mapping cone of $S^5 \cup e^7 \subset F_1 \rightarrow G_2^{(5)}$ has the homotopy type of $G_2^{(8)}$. Similarly, the homotopy fibre F_2 of $G_2^{(8)} \hookrightarrow G_2$ has a CW structure given by $S^8 \cup e^{10} \cup (\text{cells in dimensions } \geq 10)$, where the cohomology generators corresponding to S^8 and e^{10} are transgressive. Thus the mapping cone of $S^8 \cup e^{10} \subset F_2 \rightarrow G_2^{(8)}$ has the homotopy type of $G_2^{(11)}$. *QED.*

Corollary 2.1.1. $1 \geq \text{Cat}(G_2^{(5)}) \geq \text{Cat}(G_2^{(3)})$, $2 \geq \text{Cat}(G_2^{(8)}) \geq \text{Cat}(G_2^{(6)})$, $3 \geq \text{Cat}(G_2^{(11)}) \geq \text{Cat}(G_2^{(9)})$ and $4 \geq \text{Cat}(G_2)$.

Let us recall the following well-known fact due to Borel.

Fact 2.2. $H^*(G_2; \mathbb{Z}/2\mathbb{Z}) \cong \mathbb{Z}/2\mathbb{Z}[x_3, x_5]/(x_3^4, x_5^2)$.

Corollary 2.2.1. $\text{wcat}(G_2^{(5)}) \geq \text{wcat}(G_2^{(3)}) \geq 1$, $\text{wcat}(G_2^{(8)}) \geq \text{wcat}(G_2^{(6)}) \geq 2$, $\text{wcat}(G_2^{(11)}) \geq \text{wcat}(G_2^{(9)}) \geq 3$ and $\text{wcat}(G_2) \geq 4$.

Corollaries 2.1.1 and 2.2.1 yield the following.

Theorem 2.3.

<i>Skeleta</i>	$G_2^{(3)}$	$G_2^{(5)}$	$G_2^{(6)}$	$G_2^{(8)}$	$G_2^{(9)}$	$G_2^{(11)}$	G_2
<i>wcat</i>	1	1	2	2	3	3	4
<i>cat</i>	1	1	2	2	3	3	4
<i>Cat</i>	1	1	2	2	3	3	4

This completes the proof of Theorem 1.1.

3. THE RING STRUCTURE OF $h^*(Sp(3))$

To show Theorem 1.2, we introduce a cohomology theory $h^*(-)$ such that $h^*(X, A) = \{X/A, \mathbb{S}[0, 2]\}$, where $\mathbb{S}[0, 2]$ is the spectrum obtained from $\mathbb{S}$ by killing all homotopy groups of dimensions bigger than 2. Then $\mathbb{S}[0, 2]$ is a ring spectrum with $\pi_*^S(\mathbb{S}[0, 2]) \cong \mathbb{Z}[\eta]/(\eta^3, 2\eta)$, where η is the Hopf element in $\pi_1^S(\mathbb{S}) = \pi_1^S(\mathbb{S}[0, 2])$. Thus h^* is an additive and multiplicative cohomology theory with $h^* = h^*(pt) \cong \mathbb{Z}[\varepsilon]/(\varepsilon^3, 2\varepsilon)$, $\deg \varepsilon = -1$, where $\varepsilon \in h^{-1} = \pi_0^S(\Sigma^{-1}\mathbb{S}) \cong \pi_1^S(\mathbb{S})$ corresponds to η .

The characteristic map of the principal $Sp(1)$ -bundle

$$Sp(1) \hookrightarrow Sp(2) \rightarrow S^7$$

is given by $\omega = \langle \iota_3, \iota_3 \rangle : S^6 \rightarrow Sp(1) \approx S^3$ the Samelson product of two copies of the identity $\iota_3 : S^3 \rightarrow S^3$, which is a generator of $\pi_6(S^3) \cong \mathbb{Z}/12\mathbb{Z}$. We state the following well-known fact (see Whitehead [20]).

Fact 3.1. Let $\mu : S^3 \times S^3 \rightarrow S^3$ be the multiplication of $Sp(1) \approx S^3$. Then we have

$$Sp(2) \simeq S^3 \cup_{\mu \circ (1 \times \omega)} S^3 \times C(S^6) = S^3 \cup_{\omega} C(S^6) \cup_{\hat{\mu} \circ [\iota_3, \omega]^r} C(S^9),$$

where $\hat{\mu} : S^3 \times S^3 \cup_{* \times \omega} \{*\} \times C(S^6) \rightarrow S^3 \cup_{\omega} C(S^6)$ is given by $\hat{\mu}|_{S^3 \times S^3} = \mu$ and $\hat{\mu}|_{S^3 \cup_{\omega} C(S^6)} = 1$ the identity and $[\iota_3, \chi_{\omega}]^r : S^9 \rightarrow S^3 \times S^3 \cup_{* \times \omega} \{*\} \times C(S^6)$ is the relative Whitehead product of the identity $\iota_3 : S^3 \rightarrow S^3$ and the characteristic map $\chi_{\omega} : (C(S^6), S^6) \rightarrow (S^3 \cup e^7, S^3)$ of the 7-cell. Thus we have $1 \geq \text{Cat}(Sp(2)^{(3)})$, $2 \geq \text{Cat}(Sp(2)^{(7)})$ and $3 \geq \text{Cat}(Sp(2))$.

Let $\nu : S^7 \rightarrow S^4$ be the Hopf element whose suspension $\nu_n = \Sigma^{n-4}\nu$ ($n \geq 4$) gives a generator of $\pi_{n+3}(S^n) \cong \mathbb{Z}/24\mathbb{Z}$ for $n \geq 5$. Then we remark that $\omega_n = \Sigma^{n-3}\omega$ ($n \geq 3$) satisfies the formula $\omega_n = 2\nu_n \in \pi_{n+3}(S^n)$ for $n \geq 5$. By Zabrodsky [21], there is a natural splitting

$$\Sigma(S^3 \times S^3 \cup \{*\} \times (S^3 \cup_\omega e^7)) \simeq \Sigma S^3 \vee \Sigma(S^3 \cup_\omega e^7) \vee \Sigma S^3 \wedge S^3.$$

Then by the definition of a relative Whitehead product, the composition of $[\iota_3, \omega]^r$ with the projections to S^3 and $S^3 \cup_\omega e^7$ are trivial and the composition with the projection to $S^3 \wedge S^3$ is given by $\iota_3 \wedge \omega$. Thus we have

$$\Sigma(\hat{\mu} \circ [\iota_3, \omega]^r) = H(\mu) \circ \Sigma(\iota_3 \wedge \omega) = \pm \nu \circ \omega_7 = 2\nu \circ \nu_7 \neq 0$$

in $\pi_{10}(S^4) \cong \mathbb{Z}/24\mathbb{Z} \langle \nu \circ \nu_7 \rangle \oplus \mathbb{Z}/2\mathbb{Z} \langle \omega_4 \circ \nu_7 \rangle$, and hence we have

$$\Sigma^2(\hat{\mu} \circ [\iota_3, \omega]^r) = \nu_5 \circ \omega_8 = 2\nu_5^2 = 0 \in \pi_{11}(S^5) \cong \mathbb{Z}/2\mathbb{Z}$$

by Proposition 5.11 of Toda [19]. The following two facts are also well-known.

Fact 3.2. *We have the following homotopy equivalences:*

$$\begin{aligned} Sp(2)/S^3 &\simeq (S^3 \times C(S^6))/(S^3 \times S^6) = S_+^3 \wedge \Sigma(S^6) = S^7 \vee S^{10}, \\ \Sigma^2 Sp(2) &\simeq \Sigma^2(S^3 \cup_\omega C(S^6)) \vee \Sigma^2 S^{10} = S^5 \cup_{\omega_5} C(S^8) \vee S^{12}. \end{aligned}$$

Fact 3.3. *The 11-skeleton $X_{3,2}^{(11)}$ of $X_{3,2} = Sp(3)/Sp(1)$ has the homotopy type of $S^7 \cup_{\nu_7} e^{11}$.*

Restricting the principal $Sp(1)$ -bundle $Sp(1) \hookrightarrow Sp(3) \xrightarrow{q} X_{3,2}$ to the subspace $X_{3,2}^{(11)} = S^7 \cup_{\nu_7} e^{11}$ of $X_{3,2}$, we obtain the subspace $q^{-1}(X_{3,2}^{(11)}) = Sp(3)^{(14)}$ of $Sp(3)$ as the total space of the principal $Sp(1)$ -bundle $Sp(1) \hookrightarrow Sp(3)^{(14)} \xrightarrow{q} \Sigma(S^6 \cup_{\nu_6} e^{10})$ with a characteristic map $\phi : S^6 \cup_{\nu_6} e^{10} \rightarrow Sp(1) \approx S^3$, which is an extension of $\omega : S^6 \rightarrow S^3$.

Proposition 3.4. *We have the following homotopy equivalences:*

$$\begin{aligned} Sp(3)^{(14)} &\simeq S^3 \cup_{\mu \circ (1 \times \phi)} S^3 \times C(S^6 \cup_{\nu_6} e^{10}) \\ &= S^3 \cup_\phi C(S^6 \cup_{\nu_6} e^{10}) \cup C(S^9 \cup_{\nu_9} e^{13}), \\ Sp(3)^{(14)}/S^3 &\simeq (S^3 \times C(S^6 \cup_{\nu_6} e^{10}))/ (S^3 \times (S^6 \cup_{\nu_6} e^{10})) \\ &= S_+^3 \wedge \Sigma(S^6 \cup_{\nu_6} e^{10}) = (S^7 \cup_{\nu_7} e^{11}) \vee (S^{10} \cup_{\nu_{10}} e^{14}), \\ Sp(n) &\simeq Sp(n-1) \cup Sp(n-1) \times C(S^{4n-2}), \end{aligned}$$

where $Sp(n-1) \subset Sp(n)^{((2n+1)n-11)}$ for $n \geq 3$, and hence

$$\begin{aligned} Sp(n)/Sp(n)^{((2n+1)n-11)} &\simeq (Sp(n-1) \times C(S^{4n-2})) / (Sp(n-1) \times S^{4n-2} \\ &\quad \cup Sp(n-1)^{((2n-1)(n-1)-11)} \times C(S^{4n-2})) \\ &= (Sp(n-1)/Sp(n-1)^{((2n-1)(n-1)-11)}) \wedge \Sigma S^{4n-2} \\ &= \cdots = (Sp(2)/\emptyset) \wedge \Sigma S^{10} \wedge \cdots \wedge \Sigma S^{4n-2} = (Sp(2)_+) \wedge S^{(2n+1)n-10} \\ &= S^{(2n+1)n-10} \vee S^{(2n+1)n-10} \wedge Sp(2) \\ &= S^{(2n+1)n-10} \vee (S^{(2n+1)n-7} \cup_{\omega_{(2n+1)n-7}} e^{(2n+1)n-3}) \vee S^{(2n+1)n}, \quad \text{for } n \geq 3. \end{aligned}$$

This yields the following result.

Proposition 3.5. *Let $\hat{\mu} : S^3 \times S^3 \cup_{* \times \phi} \{*\} \times (S^3 \cup_{\phi} C(S^6 \cup_{\nu_6} e^{10})) \rightarrow S^3 \cup_{\phi} C(S^6 \cup_{\nu_6} e^{10})$ be the map given by $\hat{\mu}|_{S^3 \times S^3} = \mu$ and $\hat{\mu}|_{S^3 \cup_{\phi} C(S^6 \cup_{\nu_6} e^{10})} = 1$ the identity. Then we have the following cone decomposition of $Sp(3)$:*

$$Sp(3) \simeq S^3 \cup_{\phi} C(S^6 \cup_{\nu_6} e^{10}) \cup_{\hat{\mu} \circ \hat{\phi}} C(S^9 \cup_{\nu_9} e^{13}) \cup C(S^{17}) \cup C(S^{20}).$$

Corollary 3.5.1. $1 \geq \text{Cat}(Sp(3)^{(3)})$, $2 \geq \text{Cat}(Sp(3)^{(7)})$, $3 \geq \text{Cat}(Sp(3)^{(14)}) \geq \text{Cat}(Sp(3)^{(11)}) \geq \text{Cat}(Sp(3)^{(10)})$, $4 \geq \text{Cat}(Sp(3)^{(18)})$ and $5 \geq \text{Cat}(Sp(3))$.

To determine the ring structures of $h^*(Sp(2))$ and $h^*(Sp(3))$, we show the following lemma.

Lemma 3.6. *Let h^* be any multiplicative generalised cohomology theory and let $Q = S^r \cup_f e^q$ for a given map $f : S^{q-1} \rightarrow S^r$ with $h^*(Q) \cong h^*\langle 1, x, y \rangle$, where x and y correspond to the generators of $h^*(S^r) \cong h^*\langle x_0 \rangle$ and $h^*(S^q) \cong h^*\langle y_0 \rangle$. Then*

$$x^2 = \pm H^h(\lambda_2(f)) \cdot y \quad \text{in } h^*(Q),$$

where $\lambda_2(f) \in \pi_q(S^{2r})$ is the Boardman-Steer Hopf invariant equal to $\Sigma h_2(f)$ the suspension of the James-Hopf invariant $h_2(f)$ (see [1]) and $H^h : \pi_q(S^{2r}) \rightarrow h^{2r}(S^q) \cong h^{2r-q}$ is the Hurewicz homomorphism given by $H^h(g) = \Sigma_*^{-q} g^*(x_0 \otimes x_0)$.

Proof. By Boardman and Steer [1], $\overline{\Delta} : Q_2 = S^r \cup_f e^q \rightarrow Q_2 \wedge Q_2$ equals the composition $(i_2 \wedge i_2) \circ \lambda_2(f) \circ q_2$, where $q_2 : Q_2 \rightarrow Q_2/S^r = S^q$ is the collapsing map and $i_2 : S^r \hookrightarrow Q_2$ is the bottom-cell inclusion. Thus we have

$$\begin{aligned} x^2 &= \overline{\Delta}^*(x \otimes x) = ((i_2 \wedge i_2) \circ \lambda_2(f) \circ q_2)^*(x \otimes x) \\ &= q_2^*(\lambda_2(f)^*(i_2^*(x) \otimes i_2^*(x))) = q_2^*(\lambda_2(f)^*(x_0 \otimes x_0)) = q_2^*(\Sigma_*^q H^h(\lambda_2(f))). \end{aligned}$$

Since $\Sigma_*^q H^h(\lambda_2(f))$ is $H^h(\lambda_2(f)) \cdot y_0 \in h^{2r}(S^q)$ up to sign, we proceed as

$$x^2 = q_2^*(\pm H^h(\lambda_2(f)) \cdot y_0) = \pm H^h(\lambda_2(f)) \cdot q_2^*(y_0) = \pm H^h(\lambda_2(f)) \cdot y.$$

This completes the proof of the lemma. QED.

Using cohomology long exact sequences derived from the cell structure of $Sp(3)$ and a direct calculation using Proposition 3.4 and Lemma 3.6, we deduce the following result for the cohomology theory h^* considered at the beginning of this section.

Theorem 3.7. *The ring structures of $h^*(Sp(2))$ and $h^*(Sp(3))$ are as follows:*

$$\begin{aligned} h^*(Sp(2)) &\cong h^*\{1, x_3, x_7, y_{10}\}, \\ h^*(Sp(3)) &\cong h^*\{1, x_3, x_7, x_{11}, y_{10}, y_{14}, y_{18}, z_{21}\} \end{aligned}$$

with ring structures given by $x_3^2 = \varepsilon \cdot x_7$, $x_7^2 = 0$, $x_{11}^2 = 0$, $x_3 x_7 = y_{10}$, $x_3 x_{11} = y_{14}$, $x_7 x_{11} = y_{18}$ and $x_3 x_7 x_{11} = z_{21}$, where ε is the non-zero element in h^{-1} .

Remark 3.8. *The two possible attaching maps $S^{10} \rightarrow S^3 \cup_{\omega} e^7$ of e^{11} discovered by Lucía Fernández-Suárez, Antonio Gómez-Tato and Daniel Tanré [2] are homotopic in $Sp(2)$. So, we can not find any effective difference in the ring structure of $h^*(Sp(3))$ by altering, as is performed in [3], the attaching map of e^{11} .*

Corollary 3.8.1. $wcat(Sp(3)^{(3)}) \geq 1$, $wcat(Sp(3)^{(7)}) \geq 2$, $wcat(Sp(3)^{(18)}) \geq wcat(Sp(3)^{(14)}) \geq wcat(Sp(3)^{(11)}) \geq wcat(Sp(3)^{(10)}) \geq 3$ and $wcat(Sp(3)) \geq 4$, together with $wcat(Sp(2)^{(3)}) \geq 1$, $wcat(Sp(2)^{(7)}) \geq 2$ and $wcat(Sp(2)) \geq 3$.

Corollary 3.8.2.

<i>Skeleta</i>	$Sp(2)^{(3)}$	$Sp(2)^{(7)}$	$Sp(2)$
wcat	1	2	3
cat	1	2	3
Cat	1	2	3

4. PROOF OF THEOREM 1.2

By Facts 3.1 and 3.2, the smash products $\wedge^4 Sp(3)$ and $\wedge^5 Sp(3)$ satisfy

$$\begin{aligned} (\wedge^4 Sp(3))^{(19)} &\simeq S^{12} \cup_{\omega_{12}} e^{16} \vee (S^{16} \vee S^{16} \vee S^{16}) \vee (S^{19} \vee S^{19} \vee S^{19} \vee S^{19}), \\ (\wedge^5 Sp(3))^{(22)} &\simeq S^{15} \cup_{\omega_{15}} e^{19} \vee (S^{19} \vee S^{19} \vee S^{19}) \vee (S^{22} \vee S^{22} \vee S^{22} \vee S^{22}). \end{aligned}$$

Then we have the following two propositions.

Proposition 4.1. *The bottom-cell inclusions $i : S^{12} \hookrightarrow \wedge^4 Sp(3)$ and $i' : S^{15} \hookrightarrow \wedge^5 Sp(3)$ induce injective homomorphisms*

$$i_* : \pi_{18}(S^{12}) \rightarrow \pi_{18}(\wedge^4 Sp(3)^{(18)}) \quad \text{and} \quad i'_* : \pi_{21}(S^{15}) \rightarrow \pi_{21}(\wedge^5 Sp(3)),$$

respectively.

Proof. We have the following two exact sequences

$$\begin{aligned} \pi_{18}(S^{15}) &\xrightarrow{\psi} \pi_{18}(S^{12}) \xrightarrow{i_*} \pi_{18}(\wedge^4 Sp(3)^{(18)}) \rightarrow \pi_{18}(S^{16} \vee S^{16} \vee S^{16} \vee S^{16}), \\ \pi_{21}(S^{18}) &\xrightarrow{\psi'} \pi_{21}(S^{15}) \xrightarrow{i'_*} \pi_{21}(\wedge^5 Sp(3)) \rightarrow \pi_{21}(S^{19} \vee S^{19} \vee S^{19} \vee S^{19} \vee S^{19}), \end{aligned}$$

where $\pi_{18}(S^{12}) \cong \pi_{21}(S^{15}) \cong \mathbb{Z}/2\mathbb{Z}\nu_{15}^2$ and ψ and ψ' are induced from $\omega_{12} = 2\nu_{12}$ and $\omega_{15} = 2\nu_{15}$. Thus ψ and ψ' are trivial, and hence i_* and i'_* are injective. *QED.*

Proposition 4.2. *The collapsing maps $q : Sp(3)^{(18)} \rightarrow Sp(3)^{(18)}/Sp(3)^{(14)} = S^{18}$ and $q' : Sp(3) \rightarrow Sp(3)/Sp(3)^{(18)} = S^{21}$ induce injective homomorphisms*

$$\begin{aligned} q^* : \pi_{18}(\wedge^4 Sp(3)^{(18)}) &\rightarrow [Sp(3)^{(18)}, \wedge^4 Sp(3)^{(18)}] \quad \text{and} \\ q'^* : \pi_{21}(\wedge^5 Sp(3)) &\rightarrow [Sp(3), \wedge^5 Sp(3)], \end{aligned}$$

respectively.

Proof. Firstly, we show that q'^* is injective: Since we have $[Sp(3), \wedge^5 Sp(3)] = [(S^{14} \cup_{\omega_{14}} e^{18}) \vee S^{21}, \wedge^5 Sp(3)] = [S^{14} \cup_{\omega_{14}} e^{18}, \wedge^5 Sp(3)] \oplus \pi_{21}(\wedge^5 Sp(3))$ by Proposition 3.4, q'^* is clearly injective.

Secondly, we show that q^* is injective: Similarly we have $[Sp(3)^{(18)}, \wedge^4 Sp(3)^{(18)}] = [S^{14} \cup_{\omega_{14}} e^{18}, \wedge^4 Sp(3)^{(18)}]$ by Proposition 3.4. Thus it is sufficient to show that $\bar{q}^* : \pi_{18}(\wedge^4 Sp(3)^{(18)}) \rightarrow [S^{14} \cup_{\omega_{14}} e^{18}, \wedge^4 Sp(3)^{(18)}]$ is injective, where $\bar{q} : S^{14} \cup_{\omega_{14}} e^{18} \rightarrow S^{18}$ is the collapsing map. In the exact sequence

$$\pi_{15}(\wedge^4 Sp(3)^{(18)}) \xrightarrow{\omega_{15}^*} \pi_{18}(\wedge^4 Sp(3)^{(18)}) \xrightarrow{\bar{q}^*} [S^{14} \cup_{\omega_{14}} e^{18}, \wedge^4 Sp(3)^{(18)}],$$

we know that $\pi_{15}(\wedge^4 Sp(3)^{(18)}) \cong \pi_{15}(S^{12} \cup_{\omega_{12}} e^{16}) = \mathbb{Z}/2\mathbb{Z}$ is generated by the composition of ν_{12} and the bottom-cell inclusion. Since $\nu_{12} \circ \omega_{15} = 0 \in \pi_{18}(S^{12})$, the homomorphism ω_{15}^* is trivial, and hence $\bar{q}^*$ is injective. *QED.*

Then the following lemma implies that $\bar{\Delta}_4$ and $\bar{\Delta}_5$ are non-trivial by Propositions 4.1 and 4.2.

Lemma 4.3. *We obtain that $\overline{\Delta}_4 = i \circ \nu_{12}^2 \circ q : Sp(3)^{(18)} \rightarrow \wedge^4 Sp(3)^{(18)}$ and that $\overline{\Delta}_5 = i' \circ \nu_{15}^2 \circ q' : Sp(3) \rightarrow \wedge^5 Sp(3)$.*

Proof. Firstly, we show that $\overline{\Delta}_4 = i \circ \nu_{12}^2 \circ q$ implies $\overline{\Delta}_5 = i' \circ \nu_{15}^2 \circ q'$. For dimensional reasons, the image of $\overline{\Delta} : Sp(3) \rightarrow Sp(3) \wedge Sp(3)$ is in $Sp(3)^{(18)} \wedge Sp(3)^{(14)} \cup S^3 \wedge Sp(3)^{(18)}$. Since $Sp(3)^{(14)}$ is of cone-length 3 by Corollary 3.5.1, the restriction of the map $(1 \wedge \overline{\Delta}_4) \circ \overline{\Delta} = \overline{\Delta}_5$ to $Sp(3)^{(18)} \wedge Sp(3)^{(14)}$ is trivial. Thus $\overline{\Delta}_5$ equals the composition

$$\overline{\Delta}_5 : Sp(3) \rightarrow S^3 \wedge Sp(3)^{(18)} \xrightarrow{1 \wedge \overline{\Delta}_4} \wedge^5 Sp(3)^{(18)} \subset \wedge^5 Sp(3).$$

Then by $\overline{\Delta}_4 = i \circ \nu_{12}^2 \circ q$, we observe that $\overline{\Delta}_5 = i' \circ (\iota_3 \wedge \nu_{12}^2) \circ q' = i' \circ \nu_{15}^2 \circ q'$.

So, we are left to show $\overline{\Delta}_4 = i \circ \nu_{12}^2 \circ q$. For dimensional reasons, the image of $\overline{\Delta} : Sp(3)^{(18)} \rightarrow Sp(3)^{(18)} \wedge Sp(3)^{(18)}$ is in $Sp(3)^{(14)} \wedge S^3 \cup Sp(3)^{(11)} \wedge Sp(3)^{(7)} \cup Sp(3)^{(7)} \wedge Sp(3)^{(11)} \cup S^3 \wedge Sp(3)^{(14)}$. Since $S^3 \cup_\phi C(S^6 \cup_{\nu_6} e^{10})$ is of cone-length 2 by Corollary 3.5.1, the restriction of $\overline{\Delta}_3 : Sp(3)^{(18)} \rightarrow \wedge^3 Sp(3)^{(18)}$ to $S^3 \cup_\phi C(S^6 \cup_{\nu_6} e^{10})$ is trivial. Hence $1 \wedge \overline{\Delta}_3 : Sp(3)^{(14)} \wedge S^3 \cup Sp(3)^{(11)} \wedge Sp(3)^{(7)} \cup Sp(3)^{(7)} \wedge Sp(3)^{(11)} \cup S^3 \wedge Sp(3)^{(14)} \rightarrow \wedge^4 Sp(3)^{(18)}$ equals the composition

$$1 \wedge \overline{\Delta}_3 : (Sp(3) \wedge Sp(3))^{(18)} \xrightarrow{\alpha} (S^3 \cup_\omega e^7) \wedge S^{10} \cup S^3 \wedge (S^{10} \cup_{\nu_{10}} e^{14}) \xrightarrow{1 \wedge \beta} \wedge^4 (S^3 \cup_\omega e^7).$$

The map $\alpha \circ \overline{\Delta} : Sp(3)^{(18)} \rightarrow (S^3 \cup_\omega e^7) \wedge S^{10} \cup S^3 \wedge (S^{10} \cup_{\nu_{10}} e^{14})$ equals the composition

$$\alpha \circ \overline{\Delta} : Sp(3)^{(18)} \rightarrow S^{14} \cup_{\omega_{14}} e^{18} \rightarrow (S^3 \cup_\omega e^7) \wedge S^{10} \cup S^3 \wedge (S^{10} \cup_{\nu_{10}} e^{14}).$$

Collapsing the subspace $S^3 \wedge (S^{10} \cup_{\nu_{10}} e^{14})$ of $(S^3 \cup_\omega e^7) \wedge S^{10} \cup S^3 \wedge (S^{10} \cup_{\nu_{10}} e^{14})$, we obtain a map

$$q' \circ \alpha \circ \overline{\Delta} : Sp(3)^{(18)} \rightarrow S^7 \wedge S^{10},$$

where $q' : (S^3 \cup_\omega e^7) \wedge S^{10} \cup S^3 \wedge (S^{10} \cup_{\nu_{10}} e^{14}) \rightarrow S^3 \wedge S^{10}$ is the collapsing map. For dimensional reasons, $q' \circ \alpha \circ \overline{\Delta}$ equals the composition:

$$q' \circ \alpha \circ \overline{\Delta} : Sp(3)^{(18)} \rightarrow Sp(3)^{(18)} / Sp(3)^{(14)} = S^{18} \xrightarrow{\gamma} S^7 \wedge S^{10}.$$

If γ were non-trivial, then γ would be $\eta_{17} : S^{18} \rightarrow S^{17}$, and hence we should have $x_7 y_{10} = \varepsilon \cdot y_{18} \neq 0$. However, from the ring structure of $h^*(Sp(3))$ given in Theorem 3.7, we know $x_7 y_{10} = 0$, and hence we obtain $\gamma = 0$. Then the image of $\alpha \circ \overline{\Delta}$ is in the subspace $S^3 \wedge (S^{10} \cup_{\nu_{10}} e^{14})$ of $(S^3 \cup_\omega e^7) \wedge S^{10} \cup S^3 \wedge (S^{10} \cup_{\nu_{10}} e^{14})$, since they are 12-connected. Hence $\overline{\Delta}_4 = (1 \wedge \overline{\Delta}_3) \circ \overline{\Delta}$ equals the composition

$$\overline{\Delta}_4 : Sp(3)^{(18)} \xrightarrow{\alpha \circ \overline{\Delta}} S^3 \wedge (S^{10} \cup_{\nu_{10}} e^{14}) \xrightarrow{1 \wedge \beta} S^3 \wedge (\wedge^3 (S^3 \cup_\omega e^7))^{(15)} \subset \wedge^4 Sp(3)^{(18)},$$

where $(\wedge^3 (S^3 \cup_\omega e^7))^{(15)}$ is given by $(S^3 \cup_\omega e^7) \wedge S^3 \wedge S^3 \cup S^3 \wedge (S^3 \cup_\omega e^7) \wedge S^3 \cup S^3 \wedge S^3 \wedge (S^3 \cup_\omega e^7)$. Collapsing the subspace $\wedge^3 S^3$ of $(\wedge^3 (S^3 \cup_\omega e^7))^{(15)}$, we obtain a map

$$q'' \circ \beta : S^{10} \cup_{\nu_{10}} e^{14} \rightarrow S^7 \wedge S^3 \wedge S^3 \cup S^3 \wedge S^7 \wedge S^3 \cup S^3 \wedge S^3 \wedge S^7,$$

where $q'' : (\wedge^3 (S^3 \cup_\omega e^7))^{(15)} \rightarrow S^7 \wedge S^3 \wedge S^3 \cup S^3 \wedge S^7 \wedge S^3 \cup S^3 \wedge S^3 \wedge S^7$ is the collapsing map. For dimensional reasons, $q'' \circ \beta$ equals the composition

$$q'' \circ \beta : S^{10} \cup_{\nu_{10}} e^{14} \rightarrow S^{14} \xrightarrow{\gamma'} S^7 \wedge S^3 \wedge S^3 \vee S^3 \wedge S^7 \wedge S^3 \vee S^3 \wedge S^3 \wedge S^7.$$

If γ' were non-trivial, then its projection to S^{13} would be $\eta_{13} : S^{14} \rightarrow S^{13}$, and hence we should have $x_3^2 x_7 = \varepsilon \cdot y_{14} \neq 0$. However, from the ring structure of

$h^*(Sp(3))$ given in Theorem 3.7, we know $x_3^2 x_7 = \varepsilon \cdot x_7^2 = 0$, and hence we obtain $\gamma' = 0$. Hence the image of β lies in the subspace $\wedge^3 S^3$ of $\wedge^3 Sp(3)^{(18)}$.

On the other hand, for dimensional reasons, $\alpha \circ \overline{\Delta}$ equals the composition

$$\alpha \circ \overline{\Delta} : Sp(3)^{(18)} \rightarrow S^{14} \cup_{\omega_{14}} e^{18} \xrightarrow{\alpha'} S^3 \wedge (S^{10} \cup_{\nu_{10}} e^{14}),$$

where the restriction $\alpha'|_{S^{14}}$ equals the composition

$$\alpha'|_{S^{14}} : S^{14} \xrightarrow{\gamma''} S^{13} \hookrightarrow S^3 \wedge (S^{10} \cup_{\nu_{10}} e^{14}).$$

If it were non-trivial, then γ'' would be $\eta_{13} : S^{14} \rightarrow S^{13}$, and hence we should have $x_3 y_{10} = \varepsilon \cdot y_{14} \neq 0$. However, from the ring structure of $h^*(Sp(3))$ given in Theorem 3.7, we know $x_3 y_{10} = x_3^2 x_7 = \varepsilon \cdot x_7^2 = 0$, and hence we obtain $\gamma'' = 0$. Hence $\alpha \circ \overline{\Delta}$ equals the composition

$$\alpha \circ \overline{\Delta} : Sp(3)^{(18)} \xrightarrow{q} S^{18} \xrightarrow{\alpha''} S^3 \wedge (S^{10} \cup_{\nu_{10}} e^{14}),$$

and hence $\overline{\Delta}_4$ equals the composition

$$\overline{\Delta}_4 : Sp(3)^{(18)} \xrightarrow{q} S^{18} \xrightarrow{\alpha''} S^3 \wedge (S^{10} \cup_{\nu_{10}} e^{14}) \xrightarrow{1 \wedge \beta} S^3 \wedge (\wedge^3 S^3) \xrightarrow{i} \wedge^4 Sp(3)^{(18)}.$$

Now, we are ready to determine $\overline{\Delta}_4$: By Theorem 3.7, we know $x_3^2 x_{11} = \varepsilon \cdot z_{18}$ and $x_3^2 = \varepsilon \cdot x_7$, hence $\alpha'' : S^{18} \rightarrow S^{13} \cup_{\nu_{13}} e^{17}$ is a co-extension of $\eta_{16} : S^{17} \rightarrow S^{16}$ on $S^{13} \cup_{\nu_{13}} e^{17}$ and $1 \wedge \beta : S^{13} \cup_{\nu_{13}} e^{17} \rightarrow S^{12}$ is an extension of $\eta_{12} : S^{13} \rightarrow S^{12}$. Thus the composition $(1 \wedge \beta) \circ \alpha''$ is an element of the Toda bracket $\{\eta_{12}, \nu_{13}, \eta_{16}\}$ which contains a single element ν_{12}^2 by Lemma 5.12 of [19], and hence $\overline{\Delta}_4 = i \circ \nu_{12}^2 \circ q$. *QED.*

Corollary 4.3.1. $wcat(Sp(3)^{(18)}) \geq 4$ and $wcat(Sp(3)) \geq 5$.

This yields the following result.

Theorem 4.4.

<i>Skeleta</i>	$Sp(3)^{(3)}$	$Sp(3)^{(7)}$	$Sp(3)^{(10)}$	$Sp(3)^{(11)}$	$Sp(3)^{(14)}$	$Sp(3)^{(18)}$	$Sp(3)$
<i>wcat</i>	1	2	3	3	3	4	5
<i>cat</i>	1	2	3	3	3	4	5
<i>Cat</i>	1	2	3	3	3	4	5

This completes the proof of Theorem 1.2.

5. PROOF OF THEOREM 1.4

We know that for $n \geq 4$,

$$\begin{aligned} Sp(n)^{(16)} &= Sp(4)^{(15)} = Sp(3)^{(14)} \cup e^{15}, \\ Sp(n)^{(19)} &= \begin{cases} Sp(4)^{(15)} \cup (e^{18} \vee e^{18}) & n = 4, \\ Sp(4)^{(15)} \cup (e^{18} \vee e^{18}) \cup e^{19} & n \geq 5, \end{cases} \\ Sp(n)^{(21)} &= Sp(n)^{(19)} \cup e^{21} \end{aligned}$$

and that $wcat(Sp(3)^{(14)}) = cat(Sp(3)^{(14)}) = Cat(Sp(3)^{(14)}) = 3$. Firstly, we show the following.

Proposition 5.1. $wcat(Sp(4)^{(15)}) = 3$.

Proof. Since $\text{cat}(Sp(2)) = 3$, it follows that $\text{wcat}(Sp(4)^{(15)}) \geq 3$ by Theorem 5.1 of [6]. Hence we are left to show $\text{wcat}(Sp(4)^{(15)}) \leq 3$: For dimensional reasons, $\overline{\Delta}_4 = (\overline{\Delta} \wedge \overline{\Delta}) \circ \overline{\Delta} : Sp(4)^{(15)} \rightarrow \wedge^4 Sp(4)^{(15)}$ equals the composition

$$\overline{\Delta}_4 : Sp(4)^{(15)} \xrightarrow{\alpha_0} Sp(4)^{(11)} \wedge Sp(4)^{(11)} \xrightarrow{\overline{\Delta} \wedge \overline{\Delta}} \wedge^4 Sp(4)^{(11)} \hookrightarrow \wedge^4 Sp(4)^{(15)},$$

for some α_0 . By Fact 3.2, $\overline{\Delta} : Sp(4)^{(11)} \rightarrow \wedge^2 Sp(4)^{(11)}$ equals the composition

$$\overline{\Delta} : Sp(4)^{(11)} \xrightarrow{\beta_0} (S^7 \vee S^{10}) \cup e^{11} \xrightarrow{\gamma_0} \wedge^2 (S^3 \cup_\omega e^7) \hookrightarrow \wedge^2 Sp(4)^{(11)},$$

for some β_0 and γ_0 . Then for dimensional reasons, $(\beta_0 \wedge \beta_0) \circ \alpha_0 : Sp(4)^{(15)} \rightarrow ((S^7 \vee S^{10}) \cup e^{11}) \wedge ((S^7 \vee S^{10}) \cup e^{11})$ and $(\gamma_0 \wedge \gamma_0)|_{S^7 \wedge S^7} : S^7 \wedge S^7 \rightarrow \wedge^4 (S^3 \cup_\omega e^7)$ are respectively equal to the compositions

$$(\beta_0 \wedge \beta_0) \circ \alpha_0 : Sp(4)^{(15)} \xrightarrow{\alpha'_0} S^7 \wedge S^7 \hookrightarrow ((S^7 \vee S^{10}) \cup e^{11}) \wedge ((S^7 \vee S^{10}) \cup e^{11}),$$

$$(\gamma_0 \wedge \gamma_0)|_{S^7 \wedge S^7} : S^7 \wedge S^7 \xrightarrow{\gamma'_0} \wedge^4 S^3 \hookrightarrow \wedge^4 (S^3 \cup_\omega e^7),$$

for some α'_0 and γ'_0 . Hence $\overline{\Delta}_4 : Sp(4)^{(15)} \rightarrow \wedge^4 Sp(4)^{(15)}$ equals the composition

$$\overline{\Delta}_4 : Sp(4)^{(15)} \xrightarrow{\alpha'_0} S^7 \wedge S^7 \xrightarrow{\gamma'_0} \wedge^4 S^3 \hookrightarrow \wedge^4 Sp(4)^{(15)},$$

where $Sp(4)^{(15)} = Sp(3)^{(14)} \cup e^{15}$. By Theorem 3.7, $x_7^2 = 0$ in $h^*(Sp(3))$, and hence α'_0 annihilates $Sp(3)^{(14)}$. Thus $\overline{\Delta}_4 : Sp(4)^{(15)} \rightarrow \wedge^4 Sp(4)^{(15)}$ equals the composition

$$\overline{\Delta}_4 : Sp(4)^{(15)} \xrightarrow{q''} S^{15} \xrightarrow{\beta'_0} S^{14} \xrightarrow{\gamma'_0} S^{12} \xrightarrow{i''} \wedge^4 Sp(4)^{(15)}$$

for some β'_0 , where $q'' : Sp(4)^{(15)} \rightarrow Sp(4)^{(15)}/Sp(4)^{(14)} = S^{15}$ is the projection and $i'' : S^{12} = S^3 \wedge S^3 \wedge S^3 \wedge S^3 \hookrightarrow \wedge^4 Sp(4)^{(15)}$ is the inclusion. Hence the non-triviality of $\overline{\Delta}_4$ implies the non-triviality of β'_0 and γ'_0 . Therefore $\overline{\Delta}_4$ should be $i'' \circ \eta_{12}^3 \circ q''$, if it were non-trivial. However, we also know from (5.5) of [19] that η_{12}^3 is $12\nu_{12} = 6\omega_{12}$ and that $i'' \circ \omega_{12}$ is trivial by Fact 3.1. Therefore, $\overline{\Delta}_4 : Sp(4)^{(15)} \rightarrow \wedge^4 Sp(4)^{(15)}$ is trivial, and hence $\text{wcat} Sp(4)^{(15)} \leq 3$. This implies that $\text{wcat} Sp(4)^{(15)} = 3$. *QED.*

Secondly, we show the following.

Proposition 5.2. $\text{wcat}(Sp(n)^{(19)}) = 4$ for $n \geq 4$.

Proof. Since $\overline{\Delta}_5 = ((1_{Sp(n)}) \wedge \overline{\Delta}_4) \circ \overline{\Delta} : Sp(n)^{(19)} \rightarrow \wedge^5 Sp(n)^{(19)}$, it equals the composition

$$\begin{aligned} \overline{\Delta}_5 : Sp(n)^{(19)} &\xrightarrow{\overline{\Delta}} Sp(n)^{(16)} \wedge Sp(n)^{(16)} = Sp(4)^{(15)} \wedge Sp(4)^{(15)} \\ &\xrightarrow{(1_{Sp(4)^{(15)}}) \wedge \overline{\Delta}_4} \wedge^5 Sp(4)^{(15)} \hookrightarrow \wedge^5 Sp(n)^{(19)}, \end{aligned}$$

which is trivial, since $\overline{\Delta}_4 : Sp(4)^{(15)} \rightarrow \wedge^4 Sp(4)^{(15)}$ is trivial by Proposition 5.1. Thus $\text{wcat}(Sp(n)^{(19)}) \leq 4$, and hence $\text{wcat}(Sp(n)^{(19)}) = 4$. *QED.*

Let $p_j : Sp(n) \rightarrow X_{n,j} = Sp(n)/Sp(n-j)$ be the projection for $j \geq 1$. Then we have the following.

Proposition 5.3. Let $q''' : Sp(n) \rightarrow Sp(n)/Sp(n)^{((2n+1)n-3)} = S^{(2n+1)n}$ be the collapsing map and $i''' : S^{(2n+1)n-6} \hookrightarrow (\wedge^5 Sp(n)) \wedge X_{n,n-3} \wedge \cdots \wedge X_{n,1}$ the inclusion. Then

$$q'''^* \circ i'''^* : \pi_{(2n+1)n}(S^{(2n+1)n-6}) \rightarrow [Sp(n), (\wedge^5 Sp(n)) \wedge X_{n,n-3} \wedge \cdots \wedge X_{n,1}]$$

is injective.

Proof. Firstly, we have the following exact sequence

$$\begin{aligned} \pi_{(2n+1)n}(S^{(2n+1)n-3}) &\xrightarrow{\psi'''} \pi_{(2n+1)n}(S^{(2n+1)n-6}) \\ &\xrightarrow{i'''} \pi_{(2n+1)n}((\wedge^5 Sp(n)) \wedge X_{n,n-3} \wedge \cdots \wedge X_{n,1}) \rightarrow \pi_{(2n+1)n}(\vee_5 S^{(2n+1)n-2}), \end{aligned}$$

where $\pi_{(2n+1)n}(S^{(2n+1)n-6}) \cong \mathbb{Z}/2\mathbb{Z}\nu_{(2n+1)n-6}^2$ and ψ''' is induced from $\omega_{(2n+1)n-6} = 2\nu_{(2n+1)n-6}$. Thus ψ''' is trivial, and hence i'''_* is injective.

Secondly, since $(\wedge^5 Sp(n)) \wedge X_{n,n-3} \wedge \cdots \wedge X_{n,1}$ is $(n(2n+1) - 11)$ -connected, we have

$$\begin{aligned} [Sp(n), (\wedge^5 Sp(n)) \wedge X_{n,n-3} \wedge \cdots \wedge X_{n,1}] \\ = [(S^{(2n+1)n-7} \cup_{\omega_{(2n+1)n-7}} e^{(2n+1)n-3}) \vee S^{(2n+1)n}, (\wedge^5 Sp(n)) \wedge X_{n,n-3} \wedge \cdots \wedge X_{n,1}] \\ = [S^{(2n+1)n-7} \cup_{\omega_{(2n+1)n-7}} e^{(2n+1)n-3}, (\wedge^5 Sp(n)) \wedge X_{n,n-3} \wedge \cdots \wedge X_{n,1}] \\ \oplus \pi_{(2n+1)n}((\wedge^5 Sp(n)) \wedge X_{n,n-3} \wedge \cdots \wedge X_{n,1}) \end{aligned}$$

by Proposition 3.4, and hence q'''^* is injective. Thus $q'''^* \circ i'''^*$ is injective. *QED.*

Then the following lemma implies that $((1_{\wedge^5 Sp(n)}) \wedge p_{n-3} \wedge \cdots \wedge p_1) \circ \overline{\Delta}_{n+2}$ is non-trivial by Proposition 5.3, and hence we obtain Theorem 1.4.

Lemma 5.4. $((1_{\wedge^5 Sp(n)}) \wedge p_{n-3} \wedge p_{n-4} \wedge \cdots \wedge p_1) \circ \overline{\Delta}_{n+2} = i''' \circ \nu_{(2n+1)n-6}^2 \circ q'''$.

Proof. We have

$$\begin{aligned} ((1_{\wedge^5 Sp(n)}) \wedge p_{n-3} \wedge \cdots \wedge p_1) \circ \overline{\Delta}_{n+2} &= (\overline{\Delta}_5 \wedge p_{n-3} \wedge \cdots \wedge p_1) \circ \overline{\Delta}_{n-2} \\ &= (\overline{\Delta}_5 \wedge (1_{\wedge^{n-3} Sp(n)})) \circ ((1_{Sp(n)}) \wedge p_{n-3} \wedge \cdots \wedge p_1) \circ \overline{\Delta}_{n-2}. \end{aligned}$$

For dimensional reasons, the image of $((1_{Sp(n)}) \wedge p_{n-3} \wedge \cdots \wedge p_1) \circ \overline{\Delta}_{n-2}$ lies in

$$Sp(n)^{(21)} \wedge S^{15} \wedge \cdots \wedge S^{4n-1} \cup Sp(n)^{(19)} \wedge X_{n,n-3} \wedge \cdots \wedge X_{n,1}.$$

From Proposition 5.2, it follows that $\overline{\Delta}_5$ annihilates $Sp(n)^{(19)}$, and hence it equals the composition

$$\overline{\Delta}_5 : Sp(n)^{(21)} \rightarrow S^{21} \xrightarrow{\delta} \wedge^5 Sp(n)^{(21)}$$

for some $\delta \in \pi_{21}(\wedge^5 Sp(3))$. Then we obtain the following diagram except for the dotted arrow using Lemma 4.3, which is commutative up to homotopy:

$$\begin{array}{ccccc} Sp(n)^{(21)} & \xrightarrow{\quad \overline{\Delta}_5 \quad} & \wedge^5 Sp(n)^{(21)} & & \\ & \searrow & \nearrow \delta & & \uparrow j \\ & S^{21} & & & \\ & \nearrow q' & \searrow \delta_0 & & \\ Sp(3) & \xrightarrow{\quad \overline{\Delta}_5 \quad} & \wedge^5 Sp(3) & & \\ & \searrow q' & \nearrow i' & & \\ & S^{21} & \xrightarrow{\nu_{15}^2} & S^{15} & \end{array}$$

Since the pair $(\wedge^5 Sp(n), \wedge^5 Sp(3))$ is 26-connected for $n \geq 4$, we can compress δ into $\wedge^5 Sp(3)$ as $\delta \sim j \circ \delta_0$. Thus we have

$$j \circ \delta_0 \circ q' \sim j \circ i' \circ \nu_{15}^2 \circ q'.$$

Now we know that $\dim Sp(3) = 21 < 26 - 1$, and hence we can drop j from the above homotopy relation and obtain

$$\delta_0 \circ q' \sim i' \circ \nu_{15}^2 \circ q'.$$

By Proposition 4.2, $q'^* : \pi_{21}(\wedge^5 Sp(3)) \rightarrow [Sp(3), \wedge^5 Sp(3)]$ is injective, and hence we obtain

$$\delta_0 \sim i' \circ \nu_{15}^2.$$

Thus $\overline{\Delta}_5$ equals the composition

$$\overline{\Delta}_5 : Sp(n)^{(21)} \rightarrow S^{21} \xrightarrow{\nu_{15}^2} S^{15} \hookrightarrow \wedge^5 Sp(n)^{(21)}.$$

Thus $((1_{\wedge^5 Sp(n)}) \wedge p_{n-3} \wedge \cdots \wedge p_1) \circ \overline{\Delta}_{n+2}$ equals the composition

$$\begin{aligned} ((1_{\wedge^5 Sp(n)}) \wedge p_{n-3} \wedge \cdots \wedge p_1) \circ \overline{\Delta}_{n+2} : Sp(n) &\rightarrow S^{21} \wedge S^{(2n+7)(n-3)} \\ &\xrightarrow{\nu_{(2n+1)^{n-6}}^2} S^{15} \wedge S^{(2n+7)(n-3)} \hookrightarrow (\wedge^5 Sp(n)) \wedge X_{n,n-3} \wedge \cdots \wedge X_{n,1}. \end{aligned}$$

This completes the proof of the lemma. *QED.*

REFERENCES

- [1] J.M. Boardman and B. Steer, *On Hopf invariants*, Comment. Math. Helv. **42** (1967), 180–221.
- [2] L. Fernández-Suárez, A. Gómez-Tato and D. Tanré, *Hopf-Ganea invariants and weak LS category*, Top. Appl. **115** (2001), 305–316.
- [3] L. Fernández-Suárez, A. Gómez-Tato, J. Strom and D. Tanré, *The Lusternik-Schnirelmann category of $Sp(3)$* , preprint.
- [4] T. Ganea, *Some problems on numerical homotopy invariants*, Symposium on Algebraic Topology, Lect. Notes in Math. **249**, Springer Verlag, Berlin (1971), 13–22.
- [5] J. C. Gómez-Larrañaga and F. González-Acuña, *Lusternik-Schnirelmann category of 3-manifolds*, Topology **31** (1992), 791–800.
- [6] N. Iwase, *A_∞ -method in Lusternik-Schnirelmann category*, to appear in Topology.
- [7] N. Iwase, *L - S category of a sphere-bundle over a sphere*, preprint.
- [8] I. James, *The topology of Stiefel manifolds*, Cambridge University Press, Cambridge, 1976, London Math. Soc. Lec. Notes **24**.
- [9] I. James and W. Singhof, *On the category of fiber bundles, Lie groups, and Frobenius maps*, Contemp. Math. **227** (1999), 177–189.
- [10] L. Lusternik and L. Schnirelmann, *Méthodes topologiques dans les Problèmes variationnels*, Hermann, Paris, 1934.
- [11] M. Mimura and T. Nishimoto, *On the cellular decomposition of the exceptional Lie group G_2* , Proc. Amer. Math. Soc., to appear.
- [12] L. Montejano, *A quick proof of Singhof's $\text{cat}(M \times S^1) = \text{cat}(M) + 1$ theorem*, Manuscripta Math. **42** (1983), 49–52.
- [13] Y. B. Rudyak, *On the Ganea conjecture for manifolds*, Proc. Amer. Math. Soc. **125** (1997), 2511–2512.
- [14] Y. B. Rudyak, *On category weight and its applications*, Topology **38** (1999), 37–55.
- [15] P. A. Schweizer, *Secondary cohomology operations induced by the diagonal mapping*, Topology **3** (1965), 337–355.
- [16] W. Singhof, *On the Lusternik-Schnirelmann category of Lie groups*, Math. Z. **145** (1975), 111–116.
- [17] W. Singhof, *On the Lusternik-Schnirelmann category of Lie groups II*, Math. Z. **151** (1976), 143–148.

- [18] W. Singhof, *Minimal coverings of manifolds with balls*, Manuscripta Math. **29** (1979), 385–415.
- [19] H. Toda, *Composition methods in homotopy groups of spheres*, Princeton Univ. Press, Princeton, 1962, Ann. of Math. Studies, **49**.
- [20] G. W. Whitehead, *Elements of homotopy theory*, Springer Verlag, Berlin, 1978, GTM series **61**.
- [21] A. Zabrodsky, *Hopf spaces*, North-Holland, Amsterdam, 1976, North-Holland Math. Studies, **22**.

(N. Iwase) FACULTY OF MATHEMATICS, KYUSHU UNIVERSITY, ROPPONMATSU FUKUOKA 810-8560, JAPAN.

E-mail address: iwase@math.kyushu-u.ac.jp

(M. Mimura) DEPARTMENT OF MATHEMATICS, OKAYAMA UNIVERSITY, OKAYAMA 700-8530, JAPAN.

E-mail address: mimura@math.okayama-u.ac.jp